

## A simple approach to prove the addition theorem for regular solid harmonics

N C Deb and N C Sil

Department of Theoretical Physics, Indian Association for the Cultivation of Science, Jadavpur, Calcutta-700 032, India

and

P Mandal

Department of Mathematics, Visva-Bharati University, Santiniketan-731 235, West Bengal, India

*Received 28 January 1994, accepted 28 March 1994*

**Abstract** : We present a simple approach to express  $r_{12}^L Y_{LM}(\hat{r}_{12})$ , with  $r_{12} = r_1 - r_2$  as a sum of the products of  $r_1^L Y_{L_1 M_1}(\hat{r}_1)$  and  $r_2^L Y_{L_2 M_2}(\hat{r}_2)$ .

**Keywords** : Solid harmonics, addition theorem

**PACS No.** : 02.30.-f

Many problems of theoretical physics call for the representation of the regular solid harmonics  $r_{12}^L Y_{LM}(\hat{r}_{12})$  in terms of the regular solid harmonics of the vectors  $r_1$  and  $r_2$  where  $r_{12} = r_1 - r_2$ . Such a representation is generally known as the addition theorem for regular solid harmonics. There exists quite a number of approaches [1,10] to derive this addition theorem. Among them Caola's [10] approach is the simplest one. However, his derivation is based on an identity due to Hobson [11] which is not so well known. In this note, we present an alternative derivation using only the familiar results.

We start with the well known expansion

$$e^{iK \cdot (r_1 - r_2)} = 4\pi \sum_{lm} i^l j_l(Kr_{12}) Y_{lm}^*(\hat{K}) Y_{lm}(\hat{r}_{12}). \quad (1)$$

Multiplying both sides of (1) by  $Y_{LM}(\hat{K})$  and integrating over  $d\Omega_K$  we get

$$4\pi i^L j_L(Kr_{12}) Y_{LM}(\hat{r}_{12}) = \int Y_{LM}(\hat{K}) e^{iK \cdot (r_1 - r_2)} d\Omega_K. \quad (2)$$

Using similar expansions as in (1) for  $e^{iK \cdot r_1}$  and  $e^{-iK \cdot r_2}$  we get

$$\begin{aligned} \text{R.H.S. of (2)} &= (4\pi)^2 \sum_{l_1 m_1} \sum_{l_2 m_2} i^{l_1} (-i)^{l_2} j_{l_1}(Kr) j_{l_2}(Kr_2) Y_{l_1 m_1}(\hat{r}_1) \\ &\quad \times Y_{l_2 m_2}(\hat{r}_2) \int Y_{l_1 m_1}^*(\hat{K}) Y_{l_2 m_2}^*(\hat{K}) Y_{LM}(\hat{K}) d\Omega_K. \end{aligned} \quad (3)$$

The angular integration in (3) can be expressed [12] in terms of the Wigner  $3j$  symbols. Thus eq. (2) becomes

$$\begin{aligned} i^L j_L(Kr_{12}) Y_{LM}(\hat{r}_{12}) &= 4\pi \sum_{l_1 m_1} \sum_{l_2 m_2} i^{l_1} (-i)^{l_2} j_{l_1}(Kr_1) j_{l_2}(Kr_2) Y_{l_1 m_1}(\hat{r}_1) \\ &\quad \times Y_{l_2 m_2}(\hat{r}_2) C(l_1, l_2, L, m_1, m_2) \begin{pmatrix} l_1 & l_2 & L \\ -m_1 & -m_2 & M \end{pmatrix} \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (4)$$

where  $C(l_1, l_2, L, m_1, m_2)$  is a multiplying factor.

From the well known properties of the  $3j$  symbols we have

$$l_1 + l_2 \geq L \quad (5)$$

and

$$m_1 + m_2 = M,$$

where  $l_1$  and  $l_2$  are either zero or positive integers and  $-l_i \leq m_i \leq l_i$  with  $i = 1, 2$ . Now the L.H.S. of the eq. (4) contains a single spherical Bessel function of argument  $Kr_{12}$  while its R.H.S. involves product of spherical Bessel functions with arguments  $Kr_1$  and  $Kr_2$ . Using the series expansion [13]

$$j_l(Kr) = \frac{\sqrt{\pi}}{2} \sum_{s=0}^{\infty} \frac{(-1)^s (Kr/2)^{l+2s}}{s! \Gamma(l+s+3/2)} \quad (6)$$

for all those Bessel functions in (4) we find that the lowest power of  $K$  on the L.H.S. is  $L$  and this power of  $K$  will come from the various terms of the summations on the R.H.S. satisfying the relation

$$l_1 + l_2 + 2(u + v) = L \quad (7)$$

where  $u$  and  $v$  coming from the series expansions of  $j_l$ 's on the R.H.S. of (4) can have values zero or positive integers only. Eq. (7) clearly implies that

$$L \geq l_1 + l_2. \quad (8)$$

The inequalities (5) and (8) can be valid simultaneously if and only if

$$l_1 + l_2 = L. \quad (9)$$

This is the crucial point of our approach. The two double infinite summation over  $l_1 m_1$  and  $l_2 m_2$  in (4) is therefore reduced to just one *double but finite* summation over  $l_1 m_1$  (say) with  $l_1$  varying from 0 to  $L$  and  $l_2 = L - l_1$ . We then equate the coefficients of  $K^L$  from both sides of (4), noting that only  $u = v = 0$  terms need to be considered to obtain

$$\begin{aligned} 4\pi i^L r_{12}^L Y_{LM}(\hat{r}_{12}) \frac{\pi}{2^{L+1} \Gamma(L+3/2)} &= (4\pi)^2 i^L \sum_{l_1 m_1} (-1)^{l_2} r_1^{l_1} Y_{l_1 m_1}(\hat{r}_1) \\ &\times r_2^{l_2} Y_{l_2 m_2}(\hat{r}_2) \frac{\sqrt{\pi}}{2^{l_1+1} \Gamma(l_1+3/2)} \frac{\sqrt{\pi}}{2^{l_2+1} \Gamma(l_2+3/2)} \\ &\quad \begin{pmatrix} l_1 & l_2 & L \\ -m_1 & -m_2 & M \end{pmatrix} \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix} \\ &\times C(l_1, l_2, L, m_1, m_2) \end{aligned} \quad (10)$$

Under the condition (9), the  $3j$  symbols in (10) now get much simplified. The factor  $C(l_1, l_2, L, m_1, m_2)$  and the  $3j$  symbols are then obtained by comparing eqs. (107.14), (106.17) and (106.18) of reference [12] as

$$C(l_1, l_2, L, m_1, m_2) = (-1)^{m_1+m_2} (i)^{L-l_1-l_2} \left[ \frac{(2L+1)(2l_1+1)(2l_2+1)}{4\pi} \right]^{1/2}, \quad (11a)$$

$$\begin{aligned} \begin{pmatrix} l_1 & l_2 & L \\ -m_1 & -m_2 & M \end{pmatrix} &= (-1)^{l_1-l_2+m_1+m_2} \\ &\quad \frac{(2l_1)! (2l_2)! (L+M)! (L-M)!}{(2L+1)! (l_1+m_1)! (l_1-m_1)! (l_2+m_2)! (l_2-m_2)!} \quad (11b) \end{aligned}$$

and

$$\begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix} = (-1)^L \left[ \frac{(2l_1)! (2l_2)!}{(2L+1)!} \right]^{1/2} \frac{L!}{l_1! l_2!}. \quad (11c)$$

Substituting (11a)–(11c) in (10) and using the relation [14]

$$\Gamma(2z) = (2\pi)^{-1/2} 2^{2z-1/2} \Gamma(z) \Gamma(z + 1/2),$$

we finally get

$$r_{12}^L Y_{LM}(\hat{r}_{12}) = \sum_{l_1 m_1} \left[ \frac{4\pi (2L+1) (L+M)! (L-M)!}{(2l_1+1) (2l_2+1) (l_1+m_1)! (l_1-m_1)! (l_2+m_2)! (l_2-m_2)!} \right]^{1/2} \times (-1)^{l_2} r_1^{l_1} Y_{l_1 m_1}(\hat{r}_1) r_2^{l_2} Y_{l_2 m_2}(\hat{r}_2), \quad (12)$$

where

$$l_2 = L - l_1, \quad m_2 = M - m_1.$$

Eq. (12) is the required addition theorem for the regular solid harmonics.

Here we would like to mention that our approach is very similar to that of Seaton [3] only upto the eq. (3) of this article and after that his analysis is quite different.

### Acknowledgment

One of the authors (NCD) is supported by the Council of Scientific and Industrial Research, Govt. of India, New Delhi.

### References

- [1] M E Rose *J. Math. Phys.* **37** 215 (1958)
- [2] M Moshinsky *Nucl. Phys.* **13** 104 (1959)
- [3] M M J Seaton *Proc. Phys. Soc.* **77** 186 (1961)
- [4] R A Sack *J. Math. Phys.* **5** 252 (1964)
- [5] J P Dahl and M P Barnett *Molec. Phys.* **9** 175 (1965)
- [6] J D Talman *Special Functions : A Group Theoretical Approach* (New York : Benjamin) Ch.12 (1968)
- [7] K G Kay, D H Todd and H J Silverstone *J. Chem. Phys.* **51** 2359 (1969)
- [8] J M Dixon and R Lacroix *J. Phys.* **A6** 1119 (1975)
- [9] R J A Tough and A J Stone *J. Phys.* **A10** 1261 (1977)
- [10] M J Caola *J. Phys.* **A11** L23 (1978)
- [11] E W Hobson *The Theory of Spherical and Ellipsoidal Harmonics* (Cambridge : Cambridge University Press) (1931)
- [12] L D Landau and E M Lifshitz *Quantum Mechanics, Course of Theoretical Physics, Vol.3* 3rd edn (New York : Pergamon) eq. (107.14) p 442 (1985)
- [13] M Abramowitz and I A Stegun *Hand Book of Mathematical Functions* (New York : Dover) *Combination of the Formulae in Sections 10.1.1 and 9.1.10* (1970)
- [14] *Formula 6.1.18 of Reference 13*